

ACK Polyclonal Antibody

Description

Product type	Primary Antibody
Code	BT-AP00179
Host	Rabbit
Isotype	IgG
Size	20ul, 50ul, 100ul
Immunogen	The antiserum was produced against synthesized peptide derived from human ACK1. AA range:250-299
Mol wt	114327
Species reactivity	Human, Mouse
Clonality	Polyclonal
Recommended application	WB, IHC-p, ELISA
Concentration	1 mg/ml
Full name	ACK Antibody
Synonyms	TNK2; ACK1; Activated CDC42 kinase 1; ACK-1; Tyrosine kinase non-receptor protein 2

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

TNK2 encodes a tyrosine kinase that binds Cdc42Hs in its GTP-bound form and inhibits both the intrinsic and GTPase-activating protein (GAP)-stimulated GTPase activity of Cdc42Hs. This binding is mediated by a unique sequence of 47 amino acids C-terminal to an SH3 domain. The tyrosine kinase non receptor 2 may be involved in a regulatory mechanism that sustains the GTP-bound active form of Cdc43Hs and which is directly linked to a tyrosine phosphorylation signal transduction pathway. Several alternatively spliced transcript variants have been identified from this gene, but the full-length nature of only two transcript variants has been determined.


Recommended Dilution

WB: 1: 500 - 1: 2000


ELISA: 1: 10000

Not yet tested in other applications.

Images



Western Blot analysis of A549 cells using ACK Polyclonal Antibody. Secondary antibody was diluted at 1:20000



Western blot analysis of the lysates from COLO205 cells using ACK1 antibody.

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com