

CCRL1 Polyclonal Antibody

Description

| Product type | Primary Antibody |
|-------------------------|---|
| Code | BT-AP01325 |
| Host | Rabbit |
| Isotype | IgG |
| Size | 20ul, 50ul, 100ul |
| Immunogen | The antiserum was produced against synthesized peptide derived from human CCRL1. AA range:291-340 |
| Mol wt | 39914 |
| Species reactivity | Human, Mouse |
| Clonality | Polyclonal |
| Recommended application | WB, IF, ELISA |
| Concentration | 1 mg/ml |
| Full name | CCRL1 Antibody |
| Synonyms | CCRL1; CCBP2; CCR11; VSHK1; C-C chemokine receptor type 11; C-C CKR-11; CC-CKR-11; CCR-11; |
| | CC chemokine receptor-like 1; CCRL1; CCX CKR |

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


Background

The protein encoded by ACKR4 is a member of the G protein-coupled receptor family, and is a receptor for C-C type chemokines. Atypical chemokine receptor 4 has been shown to bind dendritic cell- and T cell-activated chemokines including CCL19/ELC, CCL21/SLC, and CCL25/TECK. A pseudogene of ACKR4 is found on chromosome 6. Alternatively spliced transcript variants encoding the same protein have been described.

Recommended Dilution


WB: 1: 500 - 1: 2000 IF: 1: 200 - 1: 1000 ELISA: 1: 40000 Not yet tested in other applications.

Images



Western Blot analysis of various cells using CCRL1 Polyclonal Antibody

Western blot analysis of lysates from HepG2 cells, using CCRL1 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunofluorescence analysis of MCF7 cells, using CCRL1 Antibody. The picture on the right is blocked with the synthesized peptide.

Storage -20°C for one year

> 501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com