

CD74 Polyclonal Antibody

Description

Product type Primary Antibody

Code BT-AP01540

Host Rabbit

Isotype IgG

Size 20ul, 50ul, 100ul

Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human CD74.

AA range:161-210

Mol wt 33516

Species reactivity Human

Clonality Polyclonal

Recommended application WB, IHC-p, ELISA

Concentration 1 mg/ml

Full name CD74 Antibody

Synonyms CD74; DHLAG; HLA class II histocompatibility antigen gamma chain; HLA-DR antigens-associated

invariant chain; Ia antigen-associated invariant chain; Ii; p33; CD74

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


CD74 molecule encoded by CD74 associates with class II major histocompatibility complex (MHC) and is an important chaperone that regulates antigen presentation for immune response. It also serves as cell surface receptor for the cytokine macrophage migration inhibitory factor (MIF) which, when bound to the encoded protein, initiates survival pathways and cell proliferation. This protein also interacts with amyloid precursor protein (APP) and suppresses the production of amyloid beta (Abeta). Multiple alternatively spliced transcript variants encoding different isoforms have been identified.

Recommended Dilution


WB: 1: 500 - 1: 2000 IHC-p: 1: 100 - 1: 300 ELISA: 1: 20000

Not yet tested in other applications.


Images



Western Blot analysis of PC12 cells using CD74 Polyclonal Antibody. Secondary antibody was diluted at 1:20000



 $Immun ohistochemical\ analysis\ of\ paraffin-embedded\ human-liver,\ antibody\ was\ diluted\ at\ 1:100$



Storage -20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com