

CD79b Polyclonal Antibody

Description

Product type Primary Antibody

Code BT-AP01544

Host Rabbit

Isotype IgG

Size 20ul, 50ul, 100ul

Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human

CD79B. AA range:61-110

Mol wt 26048

Species reactivity Human

Clonality Polyclonal

Recommended application WB, IHC-p, ELISA

Concentration 1 mg/ml

Full name CD79b Antibody

Synonyms CD79B; B29; IGB; B-cell antigen receptor complex-associated protein beta chain; B-cell-specific

glycoprotein B29; Ig-beta; Immunoglobulin-associated B29 protein; CD79b

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


The B lymphocyte antigen receptor is a multimeric complex that includes the antigen-specific component, surface immunoglobulin (Ig). Surface Ig non-covalently associates with two other proteins, Ig-alpha and Ig-beta, which are necessary for expression and function of the B-cell antigen receptor. CD79B encodes the Ig-beta protein of the B-cell antigen component. Alternatively spliced transcript variants encoding different isoforms have been described.

Recommended Dilution

WB: 1: 500 - 1: 2000 IHC-p: 1: 100 - 300 ELISA: 1: 20000

Not yet tested in other applications.


Images



Western Blot analysis of MCF7, L929, K562 cells using CD79b Polyclonal Antibody. Secondary antibody was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-lymph, antibody was diluted at 1:100



 $Immun ohistochemical\ analysis\ of\ paraffin-embedded\ human-lymph,\ antibody\ was\ diluted\ at\ 1:100$



 $Immun ohistochemical\ analysis\ of\ paraffin-embedded\ human-spleen,\ antibody\ was\ diluted\ at\ 1:100$



Western blot analysis of lysate from MCF7 cells, using CD79B Antibody.

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com