

Cdc37 Polyclonal Antibody

Description

Product type	Primary Antibody
Code	BT-AP01588
Host	Rabbit
Isotype	IgG
Size	20ul, 50ul, 100ul
Immunogen	The antiserum was produced against synthesized peptide derived from human p50 CDC37. AA range:231-280
Mol wt	44468
Species reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Recommended application	WB, IHC-p, IF, ELISA
Concentration	1 mg/ml
Full name	Cdc37 Antibody
Synonyms	CDC37; CDC37A; Hsp90 co-chaperone Cdc37; Hsp90 chaperone protein kinase-targeting subunit; p50Cdc37

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

Cell division cycle 37 encoded by CDC37 is highly similar to Cdc 37, a cell division cycle control protein of *Sacchomyces cerevisiae*. This protein is a molecular chaperone with specific function in cell signal transduction. It has been shown to form complex with Hsp90 and a variety of protein kinases including CDK4, CDK6, SRC, RAF-1, MOK, as well as eIF2 alpha kinases. It is thought to play a critical role in directing Hsp90 to its target kinases.

Recommended Dilution

WB: 1: 500 - 1: 2000


IHC: 1: 100 - 1: 300

IF: 1: 200 - 1: 1000


ELISA: 1: 20000

Not yet tested in other applications.

Images




Western Blot analysis of various cells using Cdc37 Polyclonal Antibody



Western blot analysis of lysates from NIH/3T3, HT-29, and RAW264.7 cells, using p50 CDC37 Antibody. The lane on the right is blocked with the synthesized peptide.

Western Blot analysis of 3T3 cells using Cdc37 Polyclonal Antibody



Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com