

Cyclin D1 Polyclonal Antibody

Description

Product type Primary Antibody

Code BT-AP02349

Host Rabbit

Isotype IgG

Size 20ul, 50ul, 100ul

Immunogen The antiserum was produced against synthesized peptide derived from human Cyclin D1. AA range:56-105

Mol wt 33729

Species reactivity Human, Mouse, Rat

Clonality Polyclonal

Recommended application WB, ELISA

Concentration 1 mg/ml

Full name Cyclin D1 Antibody

Synonyms CCND1; BCL1; PRAD1; G1/S-specific cyclin-D1; B-cell lymphoma 1 protein; BCL-1; BCL-1 oncogene;

PRAD1 oncogene

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

The protein encoded by CCND1 (cyclin D1) belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with tumor suppressor protein Rb and the expression of CCND1 is regulated positively by Rb. Mutations, amplification and overexpression of this gene, which alters cell cycle progression, are observed frequently in a variety of tumors and may contribute to tumorigenesis.


Recommended Dilution

WB: 1: 500 - 1: 2000 ELISA: 1: 10000

Not yet tested in other applications.



Cvclin D1 --



-- 34 -- 26 Western blot analysis of the lysates from K562 cells using Cyclin D1 antibody.

Western blot analysis of lysates from K562 and NIH/3T3 cells, using Cyclin D1 Antibody. The lane on the right is blocked with the synthesized peptide.

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com