

CYP2D6 Polyclonal Antibody

Description

Product type	Primary Antibody
Code	BT-AP02408
Host	Rabbit
Isotype	IgG
Size	20ul, 50ul, 100ul
Immunogen	The antiserum was produced against synthesized peptide derived from human Cytochrome P450 2D6. AA range:241-290
Mol wt	50054
Species reactivity	Human
Clonality	Polyclonal
Recommended application	WB, IHC-p, ELISA
Concentration	1 mg/ml
Full name	CYP2D6 Antibody
Synonyms	CYP2D6; CYP2DL1; Cytochrome P450 2D6; CYP11D6; Cytochrome P450-DB1; Debrisoquine 4-hydroxylase

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

CYP2D6 encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. Cytochrome P450 2D6 localizes to the endoplasmic reticulum and is known to metabolize as many as 25% of commonly prescribed drugs. Its substrates include antidepressants, antipsychotics, analgesics and antitussives, beta adrenergic blocking agents, antiarrhythmics and antiemetics. The gene is highly polymorphic in the human population; certain alleles result in the poor metabolizer phenotype, characterized by a decreased ability to metabolize the enzyme's substrates. Some individuals with the poor metabolizer phenotype have no functional protein since they carry 2 null alleles whereas in other individuals the gene is absent. CYP2D6 can vary in copy number and individuals with the ultrarapid metabolizer phenotype can have 3 or more active copies of the gene. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Recommended Dilution


WB: 1: 500 - 1: 2000

IHC: 1: 100 - 1: 300


ELISA: 1: 40000

Not yet tested in other applications.


Images




Western blot analysis of the lysates from COLO205 cells using Cytochrome P450 2D6 antibody.



Immunohistochemistry analysis of paraffin-embedded human liver, using Cytochrome P450 2D6 Antibody. The lane on the right is blocked with the Cytochrome P450 2D6 peptide.



Western blot analysis of Cytochrome P450 2D6 Antibody. The lane on the right is blocked with the Cytochrome P450 2D6 peptide.



Western Blot analysis of 293 cells using CYP2D6 Polyclonal Antibody

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com