

DDR1 Polyclonal Antibody

Description

Product type Primary Antibody

Code BT-AP02535

Host Rabbit

Isotype IgG

Size 20ul, 50ul, 100ul

Immunogen Synthesized peptide derived from DDR1 . at AA range: 740-820

Mol wt 101128

Species reactivity Human, Mouse, Rat

Clonality Polyclonal

Recommended application WB, ELISA

Concentration 1 mg/ml

Full name DDR1 Antibody

Synonyms DDR1; CAK; EDDR1; NEP; NTRK4; PTK3A; RTK6; TRKE; Epithelial discoidin domain-containing

receptor 1; Epithelial discoidin domain receptor 1; CD167 antigen-like family member A; Cell adhesion

kinase; Di

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

Receptor tyrosine kinases play a key role in the communication of cells with their microenvironment. These kinases are involved in the regulation of cell growth, differentiation and metabolism. Discoidin domain receptor tyrosine kinase 1 encoded by DDR1 belongs to a subfamily of tyrosine kinase receptors with homology to Dictyostelium discoideum protein discoidin I in their extracellular domain, and that are activated by various types of collagen. Expression of this protein is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, it has been shown to be significantly overexpressed in several human tumors. Alternatively spliced transcript variants encoding different isoforms have been described for DDR1.

Recommended Dilution

WB: 1: 500 - 1: 2000 ELISA: 1: 10000

Not yet tested in other applications.

Images

DORI 110 KD

DORI (p-Tyrs13) 110KD

+ - phospho-paptide

- - + non-phospho-paptide

Western Blot analysis of various cells using DDR1 Polyclonal Antibody diluted at 1:500. Secondary antibody was diluted at 1:20000

Storage

-20°C for one year