

ErbB-3 Polyclonal Antibody

Description

| Product type | Primary Antibody |
|-------------------------|---|
| Code | BT-AP03041 |
| Host | Rabbit |
| Isotype | IgG |
| Size | 20ul, 50ul, 100ul |
| Immunogen | Synthesized peptide derived from ErbB-3 . at AA range: 1140-1220 |
| Mol wt | 148098 |
| Species reactivity | Human, Mouse, Rat |
| Clonality | Polyclonal |
| Recommended application | WB, ELISA |
| Concentration | 1 mg/ml |
| Full name | ErbB-3 Antibody |
| Synonyms | ERBB3; HER3; Receptor tyrosine-protein kinase erbB-3; Proto-oncogene-like protein c-ErbB-3; Tyrosine kinase-type cell surface receptor HER3 |

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


Background

ERBB3 encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However, it does form heterodimers with other EGF receptor family members which do have kinase activity. Heterodimerization leads to the activation of pathways which lead to cell proliferation or differentiation. Amplification of this gene and/or overexpression of its protein have been reported in numerous cancers, including prostate, bladder, and breast tumors. Alternate transcriptional splice variants encoding different isoforms have been characterized. One isoform lacks the intermembrane region and is secreted outside the cell. This form acts to modulate the activity of the membrane-bound form. Additional splice variants have also been reported, but they have not been thoroughly characterized.

Recommended Dilution

WB: 1: 500 - 1: 2000 ELISA: 1: 40000 Not yet tested in other applications.

Images



Western Blot analysis of Jurkat cells using ErbB-3 Polyclonal Antibody cells nucleus.

Storage -20°C for one year