

Flk-1 Polyclonal Antibody

Description

Product type	Primary Antibody
Code	BT-AP03269
Host	Rabbit
Isotype	IgG
Size	20ul, 50ul, 100ul
Immunogen	The antiserum was produced against synthesized peptide derived from human VEGFR2. AA range:917-966
Mol wt	151527
Species reactivity	Human, Mouse
Clonality	Polyclonal
Recommended application	WB, IHC-p, IF, ELISA
Concentration	1 mg/ml
Full name	Flk-1 Antibody
Synonyms	KDR; FLK1; VEGFR2; Vascular endothelial growth factor receptor 2; VEGFR-2; Fetal liver kinase 1; FLK-1; Kinase insert domain receptor; KDR; Protein-tyrosine kinase receptor flk-1; CD antigen CD309

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

VEGF R1 is one of the five receptor tyrosine kinases (RTKs) (Flt-1, KDR/Flk-1, Flt-4, tie-1, and tek/tie-2) whose expression is almost exclusively restricted to the endothelial cells. Tie-1 and tek/tie-2 define a new class of RTKs containing two immunoglobulin-like domains, three EGF homology domains and three fibronectin type III domains in their extracellular regions. Flt-1, KDR/Flk-1 and Flt-4 are members of the class III subfamily of RTKs containing seven immunoglobulin-like repeats in their extracellular domains. All five RTKs are likely to play central roles in vasculogenesis and angiogenesis.

Recommended Dilution

WB: 1: 500 - 1: 2000


IHC: 1: 100 - 1: 300

IF: 1: 200 - 1: 1000


ELISA: 1: 20000

Not yet tested in other applications.


Images




Immunofluorescence analysis of rat-lung tissue. 1,Flk-1 Polyclonal Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B




Immunofluorescence analysis of rat-lung tissue. 1,Flk-1 Polyclonal Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B




Western blot analysis of lysates from SK-OV3 cells, using VEGFR2 Antibody. The lane on the right is blocked with the synthesized peptide.




Immunohistochemical analysis of paraffin-embedded Human-stomach-cancer tissue. 1,Flk-1 Polyclonal Antibody was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.




Immunofluorescence analysis of rat-kidney tissue. 1,Flk-1 Polyclonal Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B




Western Blot analysis of various cells using Flk-1 Polyclonal Antibody diluted at 1:1000




Immunofluorescence analysis of mouse-kidney tissue. 1,Flk-1 Polyclonal Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B




Immunofluorescence analysis of rat-kidney tissue. 1,Flk-1 Polyclonal Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B




Immunohistochemistry analysis of paraffin-embedded human brain tissue, using VEGFR2 Antibody. The picture on the right is blocked with the synthesized peptide.



Western Blot analysis of HeLa cells using Flk-1 Polyclonal Antibody diluted at 1:1000



Immunofluorescence analysis of mouse-kidney tissue. 1,Flk-1 Polyclonal Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B



Immunohistochemical analysis of paraffin-embedded Rat-spinal-cord tissue. 1,Flk-1 Polyclonal Antibody was diluted at 1:200(4°C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com