

## Histone H2B Polyclonal Antibody

## Description

Product type	Primary Antibody
Code	BT-AP04030
Host	Rabbit
Isotype	lgG
Size	20ul, 50ul, 100ul
Immunogen	Synthesized peptide derived from the Internal region of human Histone H2B.
Mol wt	14167/13950/13906
Species reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Recommended application	WB, IHC-p, ELISA
Concentration	1 mg/ml
Full name	Histone H2B Antibody
Synonyms	HIST1H2BB; H2BFF; Histone H2B type 1-B; Histone H2B.1; Histone H2B.f; H2B/f; HIST1H2BC; H2BFL; HIST1H2BE; H2BFH; HIST1H2BF; H2BFG; HIST1H2BG; H2BFA; HIST1H2BI; H2BFK; Histone H2B type 1-C/E/F/G/I; His

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. HIST1H2BA is intronless and encodes a replication-dependent histone that is a testis/sperm-specific member of the histone H2B family. Transcripts from HIST1H2BA contain a palindromic termination element.

## **Recommended Dilution**

WB: 1: 500 - 1: 2000 IHC-p: 1: 100 - 1: 300 ELISA: 1: 20000 Not yet tested in other applications.

Images



Immunohistochemical analysis of paraffin-embedded human-Breast-cancer, antibody was diluted at 1:100

Western Blot analysis of HeLa cells using Histone H2B Polyclonal Antibody. Secondary antibody was diluted at 1:20000 cells nucleus.

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com