

IGF-IIR Polyclonal Antibody

Description

Product type Primary Antibody

BT-AP04395 Code

Host Rabbit

Isotype IgG

20ul, 50ul, 100ul Size

The antiserum was produced against synthesized peptide derived from the C-terminal region of human Immunogen

IGF2R. AA range:2251-2300

Mol wt 274276

Species reactivity Human

Clonality Polyclonal

WB, ELISA Recommended application

Full name IGF-IIR Antibody

IGF2R; MPRI; Cation-independent mannose-6-phosphate receptor; CI Man-6-P receptor; CI-MPR; M6PR; Synonyms

300 kDa mannose 6-phosphate receptor; MPR 300;Insulin-like growth factor 2 receptor; Insulin-like growt

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

1 mg/ml

Background

Concentration


IGF2R encodes a receptor for both insulin-like growth factor 2 and mannose 6-phosphate. The binding sites for each ligand are located on different segments of the protein. This receptor has various functions, including in the intracellular trafficking of lysosomal enzymes, the activation of transforming growth factor beta, and the degradation of insulin-like growth factor 2. Mutation or loss of heterozygosity of this gene has been association with risk of hepatocellular carcinoma. The orthologous mouse gene is imprinted and shows exclusive expression from the maternal allele; however, imprinting of the human gene may be polymorphic, as only a minority of individuals showed biased expression from the maternal allele (PMID: 8267611).

Recommended Dilution

WB: 1: 500 - 1: 2000 ELISA: 1: 20000

Not yet tested in other applications.

Images



130-

Western Blot analysis of MCF7 cells using IGF-IIR Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Western blot analysis of lysate from MCF7 cells, using IGF2R Antibody.

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com