

## **KOR-1** Polyclonal Antibody

## Description

| Product type   | Primary Antibody  |
|--|---|
| Code   | BT-AP04856  |
| Host   | Rabbit  |
| Isotype  | IgG   |
| Size   | 20ul, 50ul, 100ul   |
| Immunogen  | The antiserum was produced against synthesized peptide derived from mouse KOR-1. AA range:331-380 |
| Mol wt   | 42645   |
| Species reactivity   | Mouse, Rat  |
| Clonality  | Polyclonal  |
| Recommended application  | IHC-p, WB, ELISA  |
| Concentration  | l mg/ml   |
| Full name  | KOR-1 Antibody  |
| Synonyms   | OPRK1; OPRK; Kappa-type opioid receptor; K-OR-1; KOR-1  |
| This product is for research use only not for use in human thereneutic or diagnostic presedure |   |

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


## Background

OPRK1 encodes an opioid receptor, which is a member of the 7 transmembrane-spanning G protein-coupled receptor family. It functions as a receptor for endogenous ligands, as well as a receptor for various synthetic opioids. Ligand binding results in inhibition of adenylate cyclase activity and neurotransmitter release. This opioid receptor plays a role in the perception of pain and mediating the hypolocomotor, analgesic and aversive actions of synthetic opioids. Variations in OPRK1 have also been associated with alcohol dependence and opiate addiction. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. A recent study provided evidence for translational readthrough in this gene and expression of an additional C-terminally extended isoform via the use of an alternative in-frame translation termination codon.


## **Recommended Dilution**

WB: 1: 500 - 2000 ELISA: 1: 5000 IHC: 1: 100 - 1: 300 Not yet tested in other applications.

Images



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using KOR-1 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4°over night

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com