

MAPKAPK-2 Polyclonal Antibody

Description

| Product type | Primary Antibody |
|-------------------------|--|
| Code | BT-AP05212 |
| Host | Rabbit |
| Isotype | IgG |
| Size | 20ul, 50ul, 100ul |
| Immunogen | The antiserum was produced against synthesized peptide derived from human MAPKAPK2. AA range:300- 349 |
| Mol wt | 45568 |
| Species reactivity | Human, Mouse, Rat |
| Clonality | Polyclonal |
| Recommended application | WB, IHC-p, ELISA |
| Concentration | 1 mg/ml |
| Full name | MAPKAPK-2 Antibody |
| Synonyms | MAPKAPK2; MAP kinase-activated protein kinase 2; MAPK-activated protein kinase 2; MAPKAP kinase 2; MAPKAP-K2; MAPKAPK-2; MK-2; MK2 |

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


Background

MAPKAPK2 (mitogen-activated protein kinase-activated protein kinase 2) encodes a member of the Ser/Thr protein kinase family. This kinase is regulated through direct phosphorylation by p38 MAP kinase. In conjunction with p38 MAP kinase, this kinase is known to be involved in many cellular processes including stress and inflammatory responses, nuclear export, gene expression regulation and cell proliferation. Heat shock protein HSP27 was shown to be one of the substrates of this kinase in vivo. Two transcript variants encoding two different isoforms have been found for MAPKAPK2.

Recommended Dilution

WB: 1: 500 - 1: 2000 IHC: 1: 100 - 1: 300 ELISA: 1: 20000 Not yet tested in other applications.

Images



Western blot analysis of lysates from HeLa cells, treated with Ca2+ 40nM 30', using MAPKAPK2 Antibody. The lane on the right is blocked with the synthesized peptide.

Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using MAPKAPK2 Antibody. The picture on the right is blocked with the synthesized peptide.



Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com