

MEK-4 Polyclonal Antibody

Description

Product type	Primary Antibody
Code	BT-AP05353
Host	Rabbit
Isotype	IgG
Size	20ul, 50ul, 100ul
Immunogen	The antiserum was produced against synthesized peptide derived from human SEK1/MKK4. AA range:46-95
Mol wt	44288
Species reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Recommended application	WB, ELISA
Concentration	1 mg/ml
Full name	MEK-4 Antibody
Synonyms	MAP2K4; JNKK1; MEK4; MKK4; PRKMK4; SEK1; SERK1; SKK1; Dual specificity mitogen-activated protein kinase kinase 4; MAP kinase kinase 4; MAPKK 4; JNK-activating kinase 1; MAPK/ERK kinase 4; MEK 4; SAPK/

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

MAP2K4 encodes a member of the mitogen-activated protein kinase (MAPK) family. Members of this family act as an integration point for multiple biochemical signals and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation, and development. They form a three-tiered signaling module composed of MAPKKs, MAPKKs, and MAPKs. Dual specificity mitogen-activated protein kinase kinase 4 is phosphorylated at serine and threonine residues by MAPKKs and subsequently phosphorylates downstream MAPK targets at threonine and tyrosine residues. A similar protein in mouse has been reported to play a role in liver organogenesis. A pseudogene of this gene is located on the long arm of chromosome X. Alternative splicing results in multiple transcript variants.


Recommended Dilution


WB: 1: 500 - 1: 2000

ELISA: 1: 20000

Not yet tested in other applications.

Images





Western Blot analysis of various cells using MEK-4 Polyclonal Antibody diluted at 1:1000

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com