

MOK Polyclonal Antibody

Description

| Product type | Primary Antibody |
|-------------------------|---|
| Code | BT-AP05512 |
| Host | Rabbit |
| Isotype | IgG |
| Size | 20ul, 50ul, 100ul |
| Immunogen | The antiserum was produced against synthesized peptide derived from human MOK. AA range:127-176 |
| Mol wt | 48014 |
| Species reactivity | Human, Mouse, Rat |
| Clonality | Polyclonal |
| Recommended application | IHC-p, ELISA |
| Concentration | 1 mg/ml |
| Full name | MOK Antibody |
| Synonyms | MOK; RAGE; RAGE1; MAPK/MAK/MRK overlapping kinase; MOK protein kinase; Renal tumor antigen |
| | 1; RAGE-1 |

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

MOK (MOK protein kinase) belongs to the MAP kinase superfamily. MOK was found to be regulated by caudal type transcription factor 2 (Cdx2) protein. The encoded protein, which is localized to epithelial cells in the intestinal crypt, may play a role in growth arrest and differentiation of cells of upper crypt and lower villus regions. Multiple alternatively spliced transcript variants encoding different isoforms have been observed for MOK.

Recommended Dilution

IHC: 1: 100 - 1: 300 ELISA: 1: 40000 Not yet tested in other applications.

Images

No images.

Storage -20°C for one year

-

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com