

## NMDAepsilon4 Polyclonal Antibody

### Description

<b>Product type</b>	Primary Antibody
<b>Code</b>	BT-AP06078
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Size</b>	20ul, 50ul, 100ul
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human GRIN2D. AA range:671-720
<b>Mol wt</b>	143560
<b>Species reactivity</b>	Human, Mouse, Rat, Monkey
<b>Clonality</b>	Polyclonal
<b>Recommended application</b>	WB, ELISA
<b>Concentration</b>	1 mg/ml
<b>Full name</b>	NMDAepsilon4 Antibody
<b>Synonyms</b>	GRIN2D; GluN2D; NMDAR2D; Glutamate [NMDA] receptor subunit epsilon-4; EB11; N-methyl D-aspartate receptor subtype 2D; NMDAR2D; NR2D

**This product is for research use only, not for use in human, therapeutic or diagnostic procedure.**

### Background

N-methyl-D-aspartate (NMDA) receptors are a class of ionotropic glutamate receptors. NMDA channel has been shown to be involved in long-term potentiation, an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain kinds of memory and learning. NMDA receptor channels are heteromers composed of the key receptor subunit NMDAR1 (GRIN1) and 1 or more of the 4 NMDAR2 subunits: nMDAR2A (GRIN2A), NMDAR2B (GRIN2B), NMDAR2C (GRIN2C), and NMDAR2D (GRIN2D).


### Recommended Dilution

WB: 1: 500 - 1: 2000


ELISA: 1: 40000

Not yet tested in other applications.

### Images



Western Blot analysis of 293T cells using NMDAε4 Polyclonal Antibody diluted at 1:500



Western blot analysis of lysates from COS7 cells, using GRIN2D Antibody. The lane on the right is blocked with the synthesized peptide.

## Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: [save@bt-laboratory.com](mailto:save@bt-laboratory.com) | [www.bt-laboratory.com](http://www.bt-laboratory.com)