

Olfactory receptor 10V1 Polyclonal Antibody

Description

Product type	Primary Antibody
Code	BT-AP06371
Host	Rabbit
Isotype	IgG
Size	20ul, 50ul, 100ul
Immunogen	The antiserum was produced against synthesized peptide derived from human OR10V1. AA range:161-210
Mol wt	34746
Species reactivity	Human
Clonality	Polyclonal
Recommended application	WB, IF, ELISA
Concentration	1 mg/ml
Full name	Olfactory receptor 10V1 Antibody
Synonyms	OR10V1; Olfactory receptor 10V1; Olfactory receptor OR11-256
The second set of the second second second the second second framework second	

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

Recommended Dilution


WB: 1: 500 - 1: 2000 IF: 1: 200 - 1: 1000 ELISA: 1: 20000 Not yet tested in other applications.

Images



Western Blot analysis of A549 cells using Olfactory receptor 10V1 Polyclonal Antibody

Western blot analysis of the lysates from Jurkat cells using OR10V1 antibody.



Storage -20°C for one year

> 501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com

Immunofluorescence analysis of A549 cells, using OR10V1 Antibody. The picture on the right is blocked with the synthesized peptide.

Western blot analysis of lysates from A549 and Jurkat cells, using OR10V1 Antibody. The lane on the right is blocked with the synthesized peptide.