

## Bamacan(Phospho-Ser1083) Polyclonal Antibody

### Description

|                                |  |
|--------------------------------|--|
| <b>Product type</b>            | Primary Antibody   |
| <b>Code</b>                    | BT-AP06679   |
| <b>Host</b>                    | Rabbit   |
| <b>Isotype</b>                 | IgG  |
| <b>Size</b>                    | 100ul, 50ul, 20ul  |
| <b>Immunogen</b>               | Synthesized peptide derived from human Bamacan (Phospho-Ser1083)   |
| <b>Mol wt</b>                  | N/A  |
| <b>Species reactivity</b>      | Human, Mouse, Rat  |
| <b>Clonality</b>               | Polyclonal   |
| <b>Recommended application</b> | IHC-p, IF, WB  |
| <b>Concentration</b>           | 1 mg/ml  |
| <b>Full name</b>               | Bamacan  |
| <b>Synonyms</b>                | Bamacan ;Phospho-Ser1083; Structural maintenance of chromosomes protein 3; SMC protein 3; SMC-3; Basement membrane-associated chondroitin proteoglycan; Bamacan; Chondroitin sulfate proteoglycan 6; Chromosome-associated polypeptide; hCAP |

**This product is for research use only, not for use in human, therapeutic or diagnostic procedure.**

### Background

This gene belongs to the SMC3 subfamily of SMC proteins. The encoded protein occurs in certain cell types as either an intracellular, nuclear protein or a secreted protein. The nuclear form, known as structural maintenance of chromosomes 3, is a component of the multimeric cohesin complex that holds together sister chromatids during mitosis, enabling proper chromosome segregation. Post-translational modification of the encoded protein by the addition of chondroitin sulfate chains gives rise to the secreted proteoglycan bamacan, an abundant basement membrane protein.

### Recommended Dilution

WB: 1: 500 - 1: 2000

IHC-p: 1: 50 - 1: 200

Not yet tested in other applications.

### Images

No images.

### Storage

-20°C for 1 year