

## p57 (Acetyl Lys278) Polyclonal Antibody

### Description

<b>Product type</b>	Primary Antibody
<b>Code</b>	BT-AP06841
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Size</b>	20ul, 50ul, 100ul
<b>Immunogen</b>	The antiserum was produced against synthesized Acetyl-peptide derived from human p57Kip2 around the Acetylation site of Lys278. AA range:241-290
<b>Mol wt</b>	32177
<b>Species reactivity</b>	Human, Mouse, Rat
<b>Clonality</b>	Polyclonal
<b>Recommended application</b>	WB, ELISA
<b>Concentration</b>	1 mg/ml
<b>Full name</b>	p57 (Acetyl Lys278) Antibody
<b>Synonyms</b>	CDKN1C; KIP2; Cyclin-dependent kinase inhibitor 1C; Cyclin-dependent kinase inhibitor p57; p57Kip2

**This product is for research use only, not for use in human, therapeutic or diagnostic procedure.**

### Background

CDKN1C is imprinted, with preferential expression of the maternal allele. Cyclin dependent kinase inhibitor 1C is a tight-binding, strong inhibitor of several G1 cyclin/Cdk complexes and a negative regulator of cell proliferation. Mutations in CDKN1C are implicated in sporadic cancers and Beckwith-Wiedemann syndrome, suggesting that CDKN1C is a tumor suppressor candidate. Three transcript variants encoding two different isoforms have been found for CDKN1C.


### Recommended Dilution

WB: 1: 500 - 1: 2000

ELISA: 1: 20000

Not yet tested in other applications.

### Images



Western Blot analysis of 4T1 cells using Acetyl-p57 (K278) Polyclonal Antibody. Secondary antibody was diluted at 1:20000



Western blot analysis of lysate from 4T1 cells, using p57Kip2 (Acetyl-Lys278) Antibody.

## Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: [save@bt-laboratory.com](mailto:save@bt-laboratory.com) | [www.bt-laboratory.com](http://www.bt-laboratory.com)