

# PLC Gamma1 Polyclonal Antibody

## Description

Product type Primary Antibody

Code BT-AP07262

**Host** Rabbit

Isotype IgG

**Size** 20ul, 50ul, 100ul

Immunogen The antiserum was produced against synthesized peptide derived from human PLCG1. AA range:736-785

Mol wt 148532

Species reactivity Human, Mouse, Rat, Monkey

Clonality Polyclonal

Recommended application WB, IHC-p, ELISA

Concentration 1 mg/ml

Full name PLC gamma1 Antibody

Synonyms PLCG1; PLC1; 1-phosphatidylinositol 4; 5-bisphosphate phosphodiesterase gamma-1; PLC-148;

Phosphoinositide phospholipase C-gamma-1; Phospholipase C-II; PLC-II; Phospholipase C-gamma-1; PLC-

gamma-1

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

#### Background


The protein encoded by PLCG1 (phospholipase C gamma 1) catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for this gene.

#### **Recommended Dilution**

WB: 1: 500 - 1: 2000 IHC: 1: 100 - 1: 300 ELISA: 1: 10000


Not yet tested in other applications.

## **Images**



Western Blot analysis of various cells using PLC γ1 Polyclonal Antibody diluted at 1:2000

Western blot analysis of lysates from HepG2, using PLCG1 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using PLCG1 Antibody. The picture on the right is blocked with the synthesized peptide.

# Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com