

Cdc25A(Phospho Ser178) Polyclonal Antibody

Description

Product type Primary Antibody

Code BT-AP07487

Host Rabbit

Isotype IgG

Size 20ul, 50ul, 100ul

Immunogen The antiserum was produced against synthesized peptide derived from human CDC25A around the

phosphorylation site of Ser178. AA range:144-193

Mol wt 59087

Species reactivity Human, Mouse, Rat

Clonality Polyclonal

Recommended application WB, IHC-p, IF, ELISA

Concentration 1 mg/ml

Full name M-phase inducer phosphatase 1

Synonyms M-phase inducer phosphatase 1; CDC25A; M-phase inducer phosphatase 1; Dual specificity phosphatase

Cdc25A

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


CDC25A is a member of the CDC25 family of phosphatases. CDC25A is required for progression from G1 to the S phase of the cell cycle. It activates the cyclin-dependent kinase CDC2 by removing two phosphate groups. CDC25A is specifically degraded in response to DNA damage, which prevents cells with chromosomal abnormalities from progressing through cell division. CDC25A is an oncogene, although its exact role in oncogenesis has not been demonstrated. Two transcript variants encoding different isoforms have been found for this gene.

Recommended Dilution


WB: 1: 500 - 1: 2000 IHC-p: 1: 100 - 1: 300 ELISA: 1: 10000

Not yet tested in other applications.

Images



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using CDC25A (Phospho-Ser178) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from A549 cells treated with UV 15', using CDC25A (Phospho-Ser178) Antibody. The lane on the right is blocked with the phospho peptide.

Storage

-20°C for 1 year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com