

Crystallin-alphaB(Phospho Ser45) Polyclonal Antibody

Description

Product type Primary Antibody

Code BT-AP08135

Host Rabbit

Isotype IgG

Size 20ul, 50ul, 100ul

Immunogen The antiserum was produced against synthesized peptide derived from human CRYAB around the

phosphorylation site of Ser45. AA range:21-70

Mol wt 20159

Species reactivity Human, Mouse, Rat, Monkey

Clonality Polyclonal

Recommended application WB, IHC-p, IF, ELISA

Concentration 1 mg/ml

Full name Alpha-crystallin B chain

Synonyms Alpha-crystallin B chain; CRYAB; CRYA2; Alpha-crystallin B chain; Alpha;B-crystallin; Heat shock

protein beta-5; HspB5; Renal carcinoma antigen NY-REN-27; Rosenthal fiber component

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


Mammalian lens crystallins are divided into alpha, beta, and gamma families. Alpha crystallins are composed of two gene products: alpha-A and alpha-B, for acidic and basic, respectively. Alpha crystallins can be induced by heat shock and are members of the small heat shock protein (HSP20) family. They act as molecular chaperones although they do not renature proteins and release them in the fashion of a true chaperone; instead they hold them in large soluble aggregates. Post-translational modifications decrease the ability to chaperone. These heterogeneous aggregates consist of 30-40 subunits; the alpha-A and alpha-B subunits have a 3:1 ratio, respectively. Two additional functions of alpha crystallins are an autokinase activity and participation in the intracellular architecture. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distin

Recommended Dilution

WB: 1: 500 - 1: 2000 IHC-p: 1: 100 - 1: 300 ELISA: 1: 10000

Not yet tested in other applications.

Images



Immunohistochemistry analysis of paraffin-embedded human heart, using CRYAB (Phospho-Ser45) Antibody. The picture on the right is blocked with the phospho peptide.

	117 85	Western blot analysis of lysates from COS7 cells treated with anisomycin 25ug/ml 30', using CRYAB (Phospho-Ser45) Antibody. The lane on the right is blocked with the phospho peptide.
	48	
CRYAB/Crystallin-a-B (pSer45)	34	
	26	
	19 (kD)	

Storage

-20°C for 1 year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com