

# SRF Polyclonal Antibody

## Description

| Product type            | Primary Antibody   |
|-------------------------|--|
| Code                    | BT-AP08560   |
| Host                    | Rabbit   |
| Isotype                 | IgG  |
| Size                    | 20ul, 50ul, 100ul  |
| Immunogen               | The antiserum was produced against synthesized peptide derived from human SRF. AA range:71-120 |
| Mol wt                  | 51593  |
| Species reactivity      | Human, Mouse, Rat  |
| Clonality               | Polyclonal   |
| Recommended application | WB, IHC-p, ELISA   |
| Concentration           | 1 mg/ml  |
| Full name               | SRF Antibody   |
| Synonyms                | SRF; Serum response factor; SRF  |
|                         |  |

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


# Background

SRF encodes a ubiquitous nuclear protein that stimulates both cell proliferation and differentiation. It is a member of the MADS (MCM1, Agamous, Deficiens, and SRF) box superfamily of transcription factors. Serum response factor binds to the serum response element (SRE) in the promoter region of target genes. This protein regulates the activity of many immediate-early genes, for example c-fos, and thereby participates in cell cycle regulation, apoptosis, cell growth, and cell differentiation. This gene is the downstream target of many pathways; for example, the mitogen-activated protein kinase pathway (MAPK) that acts through the ternary complex factors (TCFs). Two transcript variants encoding different isoforms have been found for this gene.

#### **Recommended Dilution**


WB: 1: 500 - 1: 2000 IHC: 1: 100 - 1: 300 ELISA: 1: 5000 Not yet tested in other applications.

#### Images



Western Blot analysis of various cells using SRF Polyclonal Antibody

Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using SRF Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from NIH/3T3 cells, treated with PMA 125ng/ml 30', using SRF Antibody. The lane on the right is blocked with the synthesized peptide.

### Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com