

TFIIIA Polyclonal Antibody

Description

Primary Antibody
BT-AP08956
Rabbit
IgG
20ul, 50ul, 100ul
Synthesized peptide derived from the Internal region of human TFIIIA.
41515
Human
Polyclonal
WB, ELISA
1 mg/ml
TFIIIA Antibody
GTF3A; Transcription factor IIIA; TFIIIA

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


Background

The product of GTF3A is a zinc finger protein with nine Cis[2]-His[2] zinc finger domains. It functions as an RNA polymerase III transcription factor to induce transcription of the 5S rRNA genes. The protein binds to a 50 bp internal promoter in the 5S genes called the internal control region (ICR), and nucleates formation of a stable preinitiation complex. This complex recruits the TFIIIC and TFIIIB transcription factors and RNA polymerase III to form the complete transcription complex. The protein is thought to be translated using a non-AUG translation initiation site in mammals based on sequence analysis, protein homology, and the size of the purified protein.

Recommended Dilution

WB: 1: 500 - 1: 2000 ELISA: 1: 40000 Not yet tested in other applications.

Images



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4° over night

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com