

## TH Polyclonal Antibody

### Description

<b>Product type</b>	Primary Antibody
<b>Code</b>	BT-AP08990
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Size</b>	20ul, 50ul, 100ul
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human Tyrosine Hydroxylase. AA range:5-54
<b>Mol wt</b>	58524
<b>Species reactivity</b>	Rat
<b>Clonality</b>	Polyclonal
<b>Recommended application</b>	WB, IHC-p, ELISA
<b>Concentration</b>	1 mg/ml
<b>Full name</b>	TH Antibody
<b>Synonyms</b>	TH; TYH; Tyrosine 3-monoxygenase; Tyrosine 3-hydroxylase; TH

**This product is for research use only, not for use in human, therapeutic or diagnostic procedure.**

### Background

Tyrosine 3-monoxygenase encoded by TH is involved in the conversion of tyrosine to dopamine. It is the rate-limiting enzyme in the synthesis of catecholamines, hence plays a key role in the physiology of adrenergic neurons. Mutations in this gene have been associated with autosomal recessive Segawa syndrome. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene.

### Recommended Dilution


WB: 1: 500 - 1: 2000

ELISA: 1: 20000


IHC-p: 1: 50 - 300

Not yet tested in other applications.


### Images



Western blot analysis of lysates from COLO205 cells, using Tyrosine Hydroxylase Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using TH Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Tyrosine Hydroxylase Antibody. The picture on the right is blocked with the synthesized peptide.

### Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: [save@bt-laboratory.com](mailto:save@bt-laboratory.com) | [www.bt-laboratory.com](http://www.bt-laboratory.com)