

VASP Polyclonal Antibody

Description

Product type Primary Antibody

BT-AP09469 Code

Host Rabbit

Isotype IgG

20ul, 50ul, 100ul Size

The antiserum was produced against synthesized peptide derived from human VASP. AA range:244-293 Immunogen

Mol wt 39699

Species reactivity Human, Mouse, Rat

Polyclonal Clonality

WB, IHC-p, ELISA Recommended application

Concentration 1 mg/ml

Full name VASP Antibody

VASP; Vasodilator-stimulated phosphoprotein; VASP Synonyms

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


Background

Vasodilator-stimulated phosphoprotein (VASP) is a member of the Ena-VASP protein family. Ena-VASP family members contain an EHV1 N-terminal domain that binds proteins containing E/DFPPPPXD/E motifs and targets Ena-VASP proteins to focal adhesions. In the midregion of the protein, family members have a proline-rich domain that binds SH3 and WW domain-containing proteins. Their C-terminal EVH2 domain mediates tetramerization and binds both G and F actin. VASP is associated with filamentous actin formation and likely plays a widespread role in cell adhesion and motility. VASP may also be involved in the intracellular signaling pathways that regulate integrinextracellular matrix interactions. VASP is regulated by the cyclic nucleotide-dependent kinases PKA and PKG.

Recommended Dilution

WB: 1: 500 - 1: 2000 IHC: 1: 100 - 1: 300 ELISA: 1: 5000

Not yet tested in other applications.



Western blot analysis of the lysates from COLO205 cells using VASP antibody.

Storage

-20°C for one year