

# **ZAP-70 Polyclonal Antibody**

## Description

Product type Primary Antibody

Code BT-AP09648

Host Rabbit

Isotype IgG

Size 20ul, 50ul, 100ul

Immunogen The antiserum was produced against synthesized peptide derived from human ZAP-70. AA range:258-307

Mol wt 69872

Species reactivity Human, Mouse

**Clonality** Polyclonal

Recommended application WB, IHC-p, ELISA

Concentration 1 mg/ml

Full name ZAP-70 Antibody

Synonyms ZAP70; SRK; Tyrosine-protein kinase ZAP-70; 70 kDa zeta-chain associated protein; Syk-related tyrosine

kinase

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

## Background

ZAP70 (zeta chain of T cell receptor associated protein kinase 70) encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in ZAP70 cause selective T-cell defect, a severe combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for ZAP70.

#### **Recommended Dilution**

WB: 1: 500 - 1: 2000 ELISA: 1: 10000


Not yet tested in other applications.

### **Images**



Western Blot analysis of 293 cells using ZAP-70 Polyclonal Antibody. Secondary antibody was

diluted at 1:20000



Western blot analysis of lysates from 293 cells, treated with PMA 125ng/ml 30', using ZAP-70 Antibody. The lane on the right is blocked with the synthesized peptide.

## Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com