

## HDAC2(Phospho Ser394) Polyclonal Antibody

Description

Product type	Primary Antibody
Code	BT-AP09847
Host	Rabbit
Isotype	IgG
Size	20ul, 50ul, 100ul
Immunogen	The antiserum was produced against synthesized peptide derived from human HDAC2 around the phosphorylation site of Ser394. AA range:360-409
Mol wt	55364
Species reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Recommended application	WB, ELISA
Concentration	1 mg/ml
Full name	Histone deacetylase 2
Synonyms	Histone deacetylase 2; HDAC2; Histone deacetylase 2; HD2


This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


## Background

This gene product belongs to the histone deacetylase family. Histone deacetylases act via the formation of large multiprotein complexes, and are responsible for the deacetylation of lysine residues at the N-terminal regions of core histones (H2A, H2B, H3 and H4). This protein forms transcriptional repressor complexes by associating with many different proteins, including YY1, a mammalian zinc-finger transcription factor. Thus, it plays an important role in transcriptional regulation, cell cycle progression and developmental events. Alternative splicing results in multiple transcript variants.

## **Recommended Dilution**

WB: 1: 500 - 1: 2000 ELISA: 1: 20000 Not yet tested in other applications.





Western blot analysis of lysates from LOVO cells, using HDAC2 (Phospho-Ser394) Antibody. The lane on the right is blocked with the phospho peptide.

Storage -20°C for 1 year

> 501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com