

HIF-1 beta Polyclonal Antibody

Description

Product type Primary Antibody

Code BT-AP09889

Host Rabbit

Isotype IgG

Size 20ul, 50ul, 100ul

Immunogen Recombinant Protein of HIF-1β

Mol wt N/A

Species reactivity Human, Mouse, Rat

Clonality Polyclonal

Recommended application WB, IHC-p, IF

Concentration

Full name Aryl hydrocarbon receptor nuclear translocator

Synonyms Aryl hydrocarbon receptor nuclear translocator; ARNT; BHLHE2; Aryl hydrocarbon receptor nuclear

translocator; ARNT protein; Class E basic helix-loop-helix protein 2; bHLHe2; Dioxin receptor, nuclear

translocator; Hypoxia-inducible factor 1-beta; HIF-1-beta; HIF1-beta

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


This gene encodes a protein containing a basic helix-loop-helix domain and two characteristic PAS domains along with a PAC domain. The encoded protein binds to ligand-bound aryl hydrocarbon receptor and aids in the movement of this complex to the nucleus, where it promotes the expression of genes involved in xenobiotic metabolism. This protein is also a co-factor for transcriptional regulation by hypoxia-inducible factor 1. Chromosomal translocation of this locus with the ETV6 (ets variant 6) gene on chromosome 12 have been described in leukemias. Alternative splicing results in multiple transcript variants.

Recommended Dilution


WB: 1: 1000 - 1: 2000 IHC: 1: 200 - 500

Not yet tested in other applications.

Images



 $Immun ohistochemical\ analysis\ of\ paraffin-embedded\ human\ Tonsil\ Tissue\ using\ HIF-1\beta\ Polyclonal\ Antibody.$



Western blot analysis of 1) Mouse Brain, 2) Rat Brain using HIF-1 β Polyclonal Antibody. Secondary antibody was diluted at 1:20000

Storage

-20°C for 1 year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com