

## MYPT1(Phospho-Ser668) Rabbit Polyclonal Antibody

### Description

<b>Product type</b>	Primary Antibody
<b>Code</b>	BT-AP11473
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Size</b>	100ul, 50ul, 20ul
<b>Immunogen</b>	Synthesized phospho peptide around human MYPT1 (Ser668)
<b>Mol wt</b>	N/A
<b>Species reactivity</b>	Human, Rat, Mouse
<b>Clonality</b>	Polyclonal
<b>Recommended application</b>	WB
<b>Concentration</b>	1 mg/ml
<b>Full name</b>	MYPT1
<b>Synonyms</b>	MYPT1 ;Ser668; Protein phosphatase 1 regulatory subunit 12A; Myosin phosphatase-targeting subunit 1; Myosin phosphatase target subunit 1; Protein phosphatase myosin-binding subunit

**This product is for research use only, not for use in human, therapeutic or diagnostic procedure.**

### Background

Myosin phosphatase target subunit 1| which is also called the myosin-binding subunit of myosin phosphatase| is one of the subunits of myosin phosphatase. Myosin phosphatase regulates the interaction of actin and myosin downstream of the guanosine triphosphatase Rho. The small guanosine triphosphatase Rho is implicated in myosin light chain (MLC) phosphorylation| which results in contraction of smooth muscle and interaction of actin and myosin in nonmuscle cells. The guanosine triphosphate (GTP)-bound| active form of RhoA (GTP.RhoA) specifically interacted with the myosin-binding subunit (MBS) of myosin phosphatase| which regulates the extent of phosphorylation of MLC. Rho-associated kinase (Rho-kinase)| which is activated by GTP. RhoA| phosphorylated MBS and consequently inactivated myosin phosphatase. Overexpression of RhoA or activated RhoA in NIH 3T3 cells increased phosph

### Recommended Dilution

WB: 1: 1000 - 1: 2000

Not yet tested in other applications.

### Images

No images.

### Storage

-20°C for 1 year