

O14AG Polyclonal Antibody

Description

Primary Antibody
BT-AP12111
Rabbit
IgG
20ul, 50ul, 100ul
Synthesized peptide derived from part region of human protein
N/A
Human, Rat, Mouse
Polyclonal
WB, ELISA
1 mg/ml
Olfactory receptor 14A16
Olfactory receptor 14A16 ;Olfactory receptor 5AT1;Olfactory receptor OR1-45

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

Recommended Dilution

WB: 1: 500 - 1: 2000 ELISA: 1: 5000 - 1: 20000 Not yet tested in other applications.

Images



Western blot analysis of lysates from HCT116 cells, primary antibody was diluted at 1:1000, 4°C overnight

Storage -20°C for 1 year

> 501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com