

## PKN3 Polyclonal Antibody

## Description

| Product type            | Primary Antibody                                                                                         |
|-------------------------|----------------------------------------------------------------------------------------------------------|
| Code                    | BT-AP13131                                                                                               |
| Host                    | Rabbit                                                                                                   |
| Isotype                 | lgG                                                                                                      |
| Size                    | 100ul, 50ul, 20ul                                                                                        |
| Immunogen               | Synthesized peptide derived from part region of human protein                                            |
| Mol wt                  | N/A                                                                                                      |
| Species reactivity      | Human, Mouse                                                                                             |
| Clonality               | Polyclonal                                                                                               |
| Recommended application | WB, ELISA                                                                                                |
| Concentration           | 1 mg/ml                                                                                                  |
| Full name               | Serine/threonine-protein kinase N3                                                                       |
| Synonyms                | Serine/threonine-protein kinase N3 ;EC 2.7.11.13;Protein kinase PKN-beta;Protein-kinase C-related kinase |
|                         | 3                                                                                                        |

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

## Background

Catalytic activity:ATP + a protein = ADP + a phosphoprotein.,The C1 domain does not bind the diacylglycerol (DAG).,enzyme regulation:Two specific sites, Thr-718 (activation loop of the kinase domain) and Thr-860 (turn motif), need to be phosphorylated for its full activation.,Contributes to invasiveness in malignant prostate cancer.,PTM:Autophosphorylated.,Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. PKC subfamily.,Contains 1 AGC-kinase C-terminal domain.,Contains 1 protein kinase domain.,Contains 3 REM (Hr1) repeats.,subcellular location:Nuclear and perinuclear Golgi region.,tissue specificity:Expressed in prostate tumors and various cancer cell lines. Not expressed in adult tissues.,

## **Recommended Dilution**

WB: 1: 500 - 1: 2000 ELISA: 1: 5000 - 1: 20000 Not yet tested in other applications.

Images No images.

Storage -20°C for 1 year

> 501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com