

U-PAR Rabbit Polyclonal Antibody

Description

Product type	Primary Antibody
Code	BT-AP13308
Host	Rabbit
Isotype	IgG
Size	100ul, 50ul, 20ul
Immunogen	Synthesized peptide derived from human U-PAR
Mol wt	36850
Species reactivity	Human, Rat, Mouse
Clonality	Polyclonal
Recommended application	WB, ELISA
Concentration	1 mg/ml
Full name	U-PAR
Synonyms	U-PAR; Urokinase plasminogen activator surface receptor; U-PAR; uPAR; Monocyte activation antigen Mo3; CD antigen CD87

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

Acts as a receptor for urokinase plasminogen activator. Plays a role in localizing and promoting plasmin formation. Mediates the proteolysis-independent signal transduction activation effects of U-PA. It is subject to negative-feedback regulation by U-PA which cleaves it into an inactive form. [Contains 3 UPAR/Ly6 domains. |subunit: Monomer (Probable). Interacts with MRC2.]

Recommended Dilution

WB: 1: 1000 - 1: 2000

ELISA: 1: 5000 - 1: 20000

Not yet tested in other applications.

Images

No images.

Storage

-20°C for 1 year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com