

PRP16 Polyclonal Antibody

Description

Product type Primary Antibody

Code BT-AP13323

Host Rabbit

Isotype IgG

Size 100ul, 50ul, 20ul

Immunogen Synthesized peptide derived from human protein . at AA range: 340-420

Mol wt N/A

Species reactivity Human, Rat, Mouse

Clonality Polyclonal

Recommended application WB, ELISA

Concentration 1 mg/ml

Full name Pre-mRNA-splicing factor ATP-dependent RNA helicase PRP16

Synonyms Pre-mRNA-splicing factor ATP-dependent RNA helicase PRP16 ;EC 3.6.4.13;ATP-dependent RNA

helicase DHX38;DEAH box protein 38

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. The protein encoded by this gene is a member of the DEAD/H box family of splicing factors. This protein resembles yeast Prp16 more closely than other DEAD/H family members. It is an ATPase and essential for the catalytic step II in pre-mRNA splicing process.

Recommended Dilution

WB: 1: 500 - 1: 2000 ELISA: 1: 5000 - 1: 20000

Not yet tested in other applications.

Images

No images.

Storage

-20°C for 1 year