

CCL14 Polyclonal Antibody

Description

Product type	Primary Antibody
Code	BT-AP14751
Host	Rabbit
Isotype	lgG
Size	20ul, 50ul, 100ul
Immunogen	Synthetic peptide from human protein at AA range: 44-93
Mol wt	N/A
Species reactivity	Human, Rat, Mouse
Clonality	Polyclonal
Recommended application	IHC-p, IF, ELISA
Concentration	1 mg/ml
Full name	C-C motif chemokine 14
Synonyms	C-C motif chemokine 14 ;Chemokine CC-1/CC-3;HCC-1/HCC-3;HCC-1;1-74;NCC-2;Small-inducible cytokine A14 [Cleaved into: HCC-1;3-74; HCC-1;4-74; HCC-1;9-74]; C-C motif chemokine 14; Chemokine CC-1/CC-3; HCC-1/HCC-3; HCC-1;1-74; NCC-2; Small-inducible cytokine A14; HCC-1;3-74; HCC-1;4-74; HCC-1;9-74;

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


Background

This gene, chemokine (C-C motif) ligand 14, is one of several CC cytokine genes clustered on 17q11.2. The CC cytokines are secreted proteins characterized by two adjacent cysteines. The cytokine encoded by this gene induces changes in intracellular calcium concentration and enzyme release in monocytes. Multiple transcript variants encoding different isoforms have been found for this gene. Read-through transcripts are also expressed that include exons from the upstream cytokine gene, chemokine (C-C motif) ligand 15, and are represented as GeneID: 348249.

Recommended Dilution

IHC-p: 1: 50 - 1: 200 ELISA: 1: 10000 - 1: 20000 Not yet tested in other applications.

Images



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200

Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com