

CD70 Polyclonal Antibody

Description

Product type	Primary Antibody
Code	BT-AP14836
Host	Rabbit
Isotype	IgG
Size	20ul, 50ul, 100ul
Immunogen	The antiserum was produced against synthesized peptide derived from human CD70. AA range:68-117
Mol wt	21118
Species reactivity	Human, Rat, Mouse
Clonality	Polyclonal
Recommended application	WB, ELISA
Concentration	1 mg/ml
Full name	CD70 antigen
Synonyms	CD70 antigen; CD70; CD27L; CD27LG; TNFSF7; CD70 antigen; CD27 ligand; CD27-L; Tumor necrosis factor ligand superfamily member 7; CD antigen CD70

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis.


Recommended Dilution

WB: 1: 500 - 1: 2000


ELISA: 1: 40000

Not yet tested in other applications.

Images



Western Blot analysis of various cells using CD70 Polyclonal Antibody diluted at 1:2000



Western blot analysis of the lysates from Jurkat cells using CD70 antibody.

Storage

-20°C for 1 year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com