

# TF(Phospho Ser290) Polyclonal Antibody

# Description

Product type Primary Antibody

Code BT-AP14873

Host Rabbit

Isotype IgG

**Size** 20ul, 50ul, 100ul

Immunogen The antiserum was produced against synthesized peptide derived from human Coagulation Factor III around

the phosphorylation site of Ser290. AA range:246-295

Mol wt 33068

Species reactivity Human, Rat, Mouse

**Clonality** Polyclonal

Recommended application WB, IHC-p, IF, ELISA

Concentration 1 mg/ml

Full name Tissue factor

Synonyms Tissue factor; F3; Tissue factor; TF; Coagulation factor III; Thromboplastin; CD antigen CD142

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

# Background


This gene encodes coagulation factor III which is a cell surface glycoprotein. This factor enables cells to initiate the blood coagulation cascades, and it functions as the high-affinity receptor for the coagulation factor VII. The resulting complex provides a catalytic event that is responsible for initiation of the coagulation protease cascades by specific limited proteolysis. Unlike the other cofactors of these protease cascades, which circulate as nonfunctional precursors, this factor is a potent initiator that is fully functional when expressed on cell surfaces. There are 3 distinct domains of this factor: extracellular, transmembrane, and cytoplasmic. This protein is the only one in the coagulation pathway for which a congenital deficiency has not been described. Alternate splicing results in multiple transcript variants.

#### Recommended Dilution


WB: 1: 500 - 1: 2000 IHC-p: 1: 100 - 1: 300 ELISA: 1: 20000

Not yet tested in other applications.

# Images



Immunohistochemistry analysis of paraffin-embedded human brain, using Coagulation Factor III (Phospho-Ser290) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from RAW264.7 cells treated with TNF 20ng/ml 30', using Coagulation Factor III (Phospho-Ser290) Antibody. The lane on the right is blocked with the phospho pentide

# Storage

-20°C for 1 year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com