

CLAP2 Polyclonal Antibody

Description

Product type	Primary Antibody
Code	BT-AP14982
Host	Rabbit
Isotype	IgG
Size	100ul, 50ul, 20ul
Immunogen	Synthesized peptide derived from part region of human protein
Mol wt	N/A
Species reactivity	Human, Rat, Mouse
Clonality	Polyclonal
Recommended application	WB, ELISA
Concentration	1 mg/ml
Full name	CLIP-associating protein 2
Synonyms	CLIP-associating protein 2 ;Cytoplasmic linker-associated protein 2;Protein Orbit homolog 2;hOrbit2

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

alternative products:Additional isoforms exist, Microtubule plus-end tracking protein that promotes the stabilization of dynamic microtubules. Required for the polarization of the cytoplasmic microtubule arrays in migrating cells towards the leading edge of the cell. May act at the cell cortex to enhance the frequency of rescue of depolymerizing microtubules by attaching their plus-ends to cortical platforms composed of ERC1 and PHLDB2. This cortical microtubule stabilizing activity is regulated at least in part by phosphatidylinositol 3-kinase signaling. Also performs a similar stabilizing function at the kinetochore which is essential for the bipolar alignment of chromosomes on the mitotic spindle. PTM:Phosphorylated by GSK3B. Phosphorylation by GSK3B may negatively regulate binding to microtubule lattices in lamella. Phosphorylated upon DNA damage, probably by ATM or ATR. Belongs to the CLASP family. Contains 5 HEAT repeats. subcellular location:Localizes to microtubule plus ends. Localizes to centrosomes, kinetochores and the mitotic spindle from prometaphase. Subsequently localizes to the spindle midzone from anaphase and to the midbody from telophase. In migrating cells localizes to the plus ends of microtubules within the cell body and to the entire microtubule lattice within the lamella. Localizes to the cell cortex and this requires ERC1 and PHLDB2. subunit:Interacts with CLIP2, ERC1, MAPRE1, MAPRE3, microtubules, PHLDB2 and RSN. The interaction with ERC1 may be mediated by PHLDB2. tissue specificity:Brain-specific.

Recommended Dilution

WB: 1: 500 - 1: 2000

ELISA: 1: 5000 - 1: 20000

Not yet tested in other applications.

Images

No images.

Storage

-20°C for 1 year