

Flt-4 Polyclonal Antibody

Description

Product type	Primary Antibody
Code	BT-AP15638
Host	Rabbit
Isotype	IgG
Size	20ul, 50ul, 100ul
Immunogen	Synthetic peptide from human protein at AA range: 720-760
Mol wt	N/A
Species reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Recommended application	IHC-p, IF, ELISA
Concentration	1 mg/ml
Full name	Vascular endothelial growth factor receptor 3
Synonyms	Vascular endothelial growth factor receptor 3 ;VEGFR-3;EC 2.7.10.1;Fms-like tyrosine kinase 4;FLT-4;Tyrosine-protein kinase receptor FLT4; Vascular endothelial growth factor receptor 3; VEGFR-3; EC 2.7.10.1; Fms-like tyrosine kinase 4; FLT-4; Tyrosine-protein kinase receptor FLT4

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

This gene encodes a tyrosine kinase receptor for vascular endothelial growth factors C and D. The protein is thought to be involved in lymphangiogenesis and maintenance of the lymphatic endothelium. Mutations in this gene cause hereditary lymphedema type IA.


Recommended Dilution

IHC-p: 1: 50 - 1: 200


ELISA: 1: 10000 - 1: 20000

Not yet tested in other applications.


Images




Immunohistochemical analysis of paraffin-embedded Human-lung-cancer, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-lung-cancer, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-placenta, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-placenta, antibody was diluted at 1:100

Storage

-20°C for 1 year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com