

ATM mouse Monoclonal Antibody(3D3)

Description

| Product type | Primary Antibody |
|-------------------------|-----------------------------------|
| Code | BT-MCA0194 |
| Host | Mouse |
| Isotype | IgG |
| Size | 20ul, 50ul, 100ul |
| Immunogen | Recombinant Protein of ATM of ATM |
| Mol wt | N/A |
| Species reactivity | Human,Rat,Mouse |
| Clonality | Monoclonal |
| Recommended application | IHC-p, IF |
| Concentration | 1 mg/ml |
| Full name | ATM |
| Synonyms | ATM |


This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

The protein encoded by this gene belongs to the PI3/PI4-kinase family. This protein is an important cell cycle checkpoint kinase that phosphorylates thus, it functions as a regulator of a wide variety of downstream proteins, including tumor suppressor proteins p53 and BRCA1, checkpoint kinase CHK2, checkpoint proteins RAD17 and RAD9, and DNA repair protein NBS1. This protein and the closely related kinase ATR are thought to be master controllers of cell cycle checkpoint signaling pathways that are required for cell response to DNA damage and for genome stability. Mutations in this gene are associated with ataxia telangiectasia, an autosomal recessive disorder.

Recommended Dilution

IHC: 1:100-200 Not yet tested in other applications.



Immunohistochemical analysis of paraffin-embedded Human Colon Carcinoma Tissue using ATM Mouse Monoclonal antibody diluted at 1:200.

Immunohistochemical analysis of paraffin-embedded Human Breast Carcinoma Tissue using ATM Mouse Monoclonal antibody diluted at 1:200.

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com