

CaMKII Beta/Gamma/Delta (Phospho Thr287) Monoclonal Antibody(1B9)

Description

Product type Primary Antibody

Code BT-MCA0252

Host Mouse

Isotype IgG

Size 20ul, 50ul, 100ul

 Immunogen
 Synthetic Peptide of CaMKIIBeta/γ/δ (Phospho Thr287)

Mol wt N/A

Species reactivity Human, Rat, Mouse

Clonality Monoclonal

Recommended application IHC-p, IF

Concentration 1 mg/ml

Full name

Synonyms CAMK2B; CAMK2; CAMK8; Calcium; calmodulin-dependent protein kinase type II subunit beta;

CaM kinase II subunit beta; CaMK-II subunit beta; CAMK2G; CAMK; CAMK-II; CAMKG; Calcium;

calmodulin-dependent protein kinase type II subunit gamma;

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


The product of this gene belongs to the serine/threonine protein kinase family and to the Ca(2+)/calmodulin-dependent protein kinase subfamily. Calcium signaling is crucial for several aspects of plasticity at glutamatergic synapses. In mammalian cells, the enzyme is composed of four different chains: alpha, beta, gamma, and delta. The product of this gene is a beta chain. It is possible that distinct isoforms of this chain have different cellular localizations and interact differently with calmodulin. Alternative splicing results in multiple transcript variants.

Recommended Dilution

IHC: 1:100-200

Not yet tested in other applications.

Images



 $Immunohistochemical\ analysis\ of\ paraffin-embedded\ Rat\ Brain\ Tissue\ using\ CaMKIIBeta/\ Gamma\ /\delta\ (Phospho\ Thr287)\ (Monoclonal\ antibody\ diluted\ at\ 1:200.$



 $Immunohistochemical\ analysis\ of\ paraffin-embedded\ Mouse\ Brain\ Tissue\ using\ CaMKIIBeta/Gamma\ /\delta\ (Phospho\ Thr 287)\ Mouse\ Monoclonal\ antibody\ diluted\ at\ 1:200.$

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com