

## CD15 Monoclonal Antibody(Q89)

### Description

<b>Product type</b>	Primary Antibody
<b>Code</b>	BT-MCA0276
<b>Host</b>	Mouse
<b>Isotype</b>	IgG
<b>Size</b>	20ul, 50ul, 100ul
<b>Immunogen</b>	Synthetic Peptide of CD15
<b>Mol wt</b>	45570
<b>Species reactivity</b>	Human
<b>Clonality</b>	Monoclonal
<b>Recommended application</b>	IHC-P, IF, ICC
<b>Concentration</b>	1 mg/ml
<b>Full name</b>	Alpha-(1,3)-fucosyltransferase 4
<b>Synonyms</b>	FUT4; ELFT; FCT3A; Alpha-(1; 3)-fucosyltransferase; ELAM-1 ligand fucosyltransferase; Fucosyltransferase 4; Fucosyltransferase IV; Fuc-TIV; FucT-IV; Galactoside 3-L-fucosyltransferase

**This product is for research use only, not for use in human, therapeutic or diagnostic procedure.**

### Background

The product of this gene transfers fucose to N-acetylglucosamine polysaccharides to generate fucosylated carbohydrate structures. It catalyzes the synthesis of the non-sialylated antigen, Lewis x (CD15).


### Recommended Dilution

IF: 1:50-200


IHC: 1:200

Not yet tested in other applications.


### Images




Immunohistochemical analysis of paraffin-embedded Human-lung-cancer tissue. 1. CD15 Monoclonal antibody(Q89) was diluted at 1:200(4°C, overnight). 2. Sodium citrate pH 6.0 was used for antibody retrieval(>98°C, 20min). 3. Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.




Immunofluorescence analysis of Human-liver-cancer tissue. 1. CD15 Monoclonal antibody(Q89)(red) was diluted at 1:200(4°C, overnight). 2. Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3. Picture B: DAPI(blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B




Immunohistochemical analysis of paraffin-embedded Rat Brain Tissue using CD 15 Mouse Monoclonal antibody diluted at 1:500.



Immunohistochemical analysis of paraffin-embedded Human stomach.1.Antibody was diluted at 1:200(4°C overnight). 2.High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3.Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human stomach.1.Antibody was diluted at 1:200(4°C overnight). 2.High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3.Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human stomach.1.Antibody was diluted at 1:200(4°C overnight). 2.High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3.Secondary antibody was diluted at 1:200(room temperature, 30min).

### Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: [save@bt-laboratory.com](mailto:save@bt-laboratory.com) | [www.bt-laboratory.com](http://www.bt-laboratory.com)