

Dynamin II Monoclonal Antibody

Description

Product type	Primary Antibody
Code	BT-MCA0451
Host	Mouse
Isotype	IgG
Size	50ul, 100ul
Immunogen	Purified recombinant fragment of Dynamin II expressed in E. Coli.
Mol wt	N/A
Species reactivity	Human
Clonality	Monoclonal
Recommended application	WB, IHC-p, IF, ELISA
Concentration	1 mg/ml
Full name	Dynamin-2
Synonyms	DNM2; DYN2; Dynamin-2

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

Dynamins represent one of the subfamilies of GTP-binding proteins. These proteins share considerable sequence similarity over the N-terminal portion of the molecule, which contains the GTPase domain. Dynamins are associated with microtubules. They have been implicated in cell processes such as endocytosis and cell motility, and in alterations of the membrane that accompany certain activities such as bone resorption by osteoclasts. Dynamins bind many proteins that bind actin and other cytoskeletal proteins. Dynamins can also self-assemble, a process that stimulates GTPase activity. Five alternatively spliced transcripts encoding different proteins have been described. Additional alternatively spliced transcripts may exist, but their full-length nature has not been determined.

Recommended Dilution


ELISA: 1:10000

IHC: 1:200 - 1:1000


WB: 1:500 - 1:2000

Not yet tested in other applications.

Images



Western Blot analysis using Dynamin II Monoclonal antibody against truncated Dynamin-2 recombinant protein (1) SKN-SH cell lysate (2) and NIH/3T3 cell lysate (3).



Immunohistochemistry analysis of paraffin-embedded human cerebrum tissue (left) and myelencephalon tissue (right), showing cytoplasmic localization with DAB staining using Dynamin II Monoclonal antibody.

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China
Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com