

Factor VIII Monoclonal Antibody

Description

| | |
|--------------------------------|--|
| Product type | Primary Antibody |
| Code | BT-MCA0524 |
| Host | Mouse |
| Isotype | IgG |
| Size | 50ul, 100ul |
| Immunogen | Purified recombinant fragment of Factor VIII expressed in E. Coli. |
| Mol wt | N/A |
| Species reactivity | Human |
| Clonality | Monoclonal |
| Recommended application | WB, ELISA |
| Concentration | 1 mg/ml |
| Full name | Coagulation factor VIII |
| Synonyms | F8; F8C; Coagulation factor VIII; Antihemophilic factor; AHF; Procoagulant component |

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

This gene encodes coagulation factor VIII, which participates in the intrinsic pathway of blood coagulation| factor VIII is a cofactor for factor IXa which, in the presence of Ca²⁺ and phospholipids, converts factor X to the activated form Xa. This gene produces two alternatively spliced transcripts. Transcript variant 1 encodes a large glycoprotein, isoform a, which circulates in plasma and associates with von Willebrand factor in a noncovalent complex. This protein undergoes multiple cleavage events. Transcript variant 2 encodes a putative small protein, isoform b, which consists primarily of the phospholipid binding domain of factor VIIIc. This binding domain is essential for coagulant activity. Defects in this gene results in hemophilia A, a common recessive X-linked coagulation disorder.


Recommended Dilution

ELISA: 1:10000

WB: 1:500 - 1:2000

Not yet tested in other applications.

Images



Western Blot analysis using Factor VIII Monoclonal antibody against truncated Trx-Factor VIII recombinant protein (1).

Storage

-20°C for one year