

Histone H3 Monoclonal Antibody(8F7)

Description

Primary Antibody
BT-MCA0712
Mouse
IgG
20ul, 50ul, 100ul
Recombinant Protein of Histone H3
15273
Human,Mouse,Rat
Monoclonal
WB
l mg/ml
Histone H3.2
HIST1H3A; H3FA; HIST1H3B; H3FL; HIST1H3C; H3FC; HIST1H3D; H3FB; HIST1H3E; H3FD; HIST1H3F; H3FI; HIST1H3G; H3FH; HIST1H3H; H3FK; HIST1H3I; H3FF; HIST1H3J; H3FJ; Histone H3.1; Histone H3; a; Histone H3; b; Histone H3; c; Histone H3; d; Histone H3; f; Histone H3; h; Histone H3; i; Histone H3; j; Histone H3; k; Histone H3; l; HIST2H3A; HIST2H3C; H3F2; H3FM; HIST2H3D; Histone H3.2; Histone H3; m; Histone H3; o; H3F3A; H3.AA; H3F3; PP781; H3F3B; H3.3B; Histone H3.3

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails| instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3.

Recommended Dilution

WB: 1:1000 -2000 Not yet tested in other applications.

Images



Western blot analysis of 1) Hela, 2) Raw 264.7, 3) Rat Testis diluted at 1:2000.

Storage -20°C for one year