

HSC70 Monoclonal Antibody(8G7)

Description

Product type Primary Antibody

Code BT-MCA0738

Host Mouse

Isotype IgG

Size 20ul, 50ul, 100ul

Immunogen Synthesized peptide derived from the Internal region of human HSC 70.

Mol wt N/A

Species reactivity Human, Rat, Mouse

Clonality Monoclonal

Recommended application WB, IHC-p, IF

Concentration 1 mg/ml

Full name Heat shock cognate 71 kDa protein (Heat shock 70 kDa protein 8)

Synonyms Heat shock cognate 71 kDa protein; Heat shock 70 kDa protein 8

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


This gene encodes a member of the heat shock protein 70 family, which contains both heat-inducible and constitutively expressed members. This protein belongs to the latter group, which are also referred to as heat-shock cognate proteins. It functions as a chaperone, and binds to nascent polypeptides to facilitate correct folding. It also functions as an ATPase in the disassembly of clathrin-coated vesicles during transport of membrane components through the cell. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Recommended Dilution


IHC-p: 1:50-300 WB: 1:500 - 1:2000

Not yet tested in other applications.

Images



Western blot analysis of 1)Hela, 2)Mouse Brain Tissue, 3) Rat Brain Tissue with HSC70 Mouse Monoclonal antibody diluted at 1:2.000.



 $Immun ohistochemical \ analysis \ of paraffin-embedded \ Human \ Breast \ Caricnoma \ using \ HSC70 \ Mouse \\ Monoclonal \ antibody \ diluted \ at \ 1:200.$

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com