

HXK I Monoclonal Antibody

Description

Product type Primary Antibody

Code BT-MCA0759

Host Mouse

Isotype IgG

Size 50ul, 100ul

Immunogen Purified recombinant fragment of human HXK I expressed in E. Coli.

Mol wt N/A

Species reactivity Human, Mouse, Rat

Clonality Monoclonal

Recommended application WB, IHC-p, IF, ICC, ELISA

Concentration 1 mg/ml

Full name Hexokinase-1

Synonyms HK1; Hexokinase-1; Brain form hexokinase; Hexokinase type I; HK I

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. This gene encodes a ubiquitous form of hexokinase which localizes to the outer membrane of mitochondria. Mutations in this gene have been associated with hemolytic anemia due to hexokinase deficiency. Alternative splicing of this gene results in several transcript variants which encode different isoforms, some of which are tissue-specific.

Recommended Dilution


ELISA: 1:10000 IF: 1:200 - 1:1000 IHC: 1:200 - 1:1000 WB: 1:500 - 1:2000

Not yet tested in other applications.


Images



Western Blot analysis using HXK I Monoclonal antibody against Jurkat (1) HeLa (2) HepG2 (3) MCF-7 (4) and PC-12 (5) cell lysate.



Immunohistochemistry analysis of paraffin-embedded human salivary gland tissues (left) and kidney tissues (right) with DAB staining using HXK I Monoclonal antibody.



Immunofluorescence analysis of NIH/3T3 cells using HXK I Monoclonal antibody (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.

Storage -20°C for one year

 $501\ Changsheng\ S\ Rd,\ Nanhu\ Dist,\ Jiaxing,\ Zhejiang,\ China$ Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com