

HXK II Monoclonal Antibody

Description

Product type	Primary Antibody
Code	BT-MCA0760
Host	Mouse
Isotype	IgG
Size	50ul, 100ul
Immunogen	Purified recombinant fragment of human HXK II expressed in E. Coli.
Mol wt	N/A
Species reactivity	Human
Clonality	Monoclonal
Recommended application	WB, IHC-p, IF, FCM, ELISA
Concentration	1 mg/ml
Full name	Hexokinase-2
Synonyms	HK2; Hexokinase-2; Hexokinase type II; HK II; Muscle form hexokinase

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. This gene encodes hexokinase 2, the predominant form found in skeletal muscle. It localizes to the outer membrane of mitochondria. Expression of this gene is insulin-responsive, and studies in rat suggest that it is involved in the increased rate of glycolysis seen in rapidly growing cancer cells.

Recommended Dilution

ELISA: 1:10000


FC: 1:200 - 1:400

IHC: 1:200 - 1:1000


WB: 1:500 - 1:2000

Not yet tested in other applications.


Images



Western Blot analysis using HXK II Monoclonal antibody against Jurkat (1) HeLa (2) and HEK293 (3) cell lysate.



Immunohistochemistry analysis of paraffin-embedded esophagus cancer tissues (left) and human lung cancer (right) with DAB staining using HXK II Monoclonal antibody.



Flow cytometric analysis of K562 cells using HXK II Monoclonal antibody (green) and negative control (purple).

Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com