

MEK2 Monoclonal Antibody(7D11)

Description

Product type	Primary Antibody
Code	BT-MCA0881
Host	Mouse
Isotype	IgG
Size	20ul, 50ul, 100ul
Immunogen	Synthesized peptide derived from human MEK-2 around the non-phosphorylation site of T394.
Mol wt	N/A
Species reactivity	Human,Rat,Mouse
Clonality	Monoclonal
Recommended application	WB
Concentration	1 mg/ml
Full name	Dual specificity mitogen-activated protein kinase kinase 2
Synonyms	MAP2K2; MEK2; MKK2; PRKMK2; Dual specificity mitogen-activated protein kinase kinase 2; MAP kinase kinase 2; MAPKK 2; ERK activator kinase 2; MAPK; ERK kinase 2; MEK 2

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background


The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. Mutations in this gene cause cardiofaciocutaneous syndrome (CFC syndrome), a disease characterized by heart defects, mental retardation, and distinctive facial features similar to those found in Noonan syndrome. The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for this gene.

Recommended Dilution

WB: 1:500 - 1:2000

Not yet tested in other applications.

Images



Western blot analysis of 1) Hela, 2)3T3. 3) Rat Brain Tissue with MEK2 Mouse Monoclonal antibody diluted at 1:2.000.

Storage

-20°C for one year